




Hi everyone, it is Felix. Dr. J. and I had a wonderful sunrise run at town lake this morning. How do you like my stylin' green LED collar that lights up so other runners see me before the sun is up?


# Special Alkene Bonus: Important material you will need to know!

Alkene stability part 1: Z (cis) groups larger than H atoms will crunch into each other causing steric strain.



Alkene stability part 2: For reasons we are not able to tell you, more substituted alkenes have more stable (stronger) pi bonds than alkenes with more H atoms on their  $sp^2$ -hybridized C atoms (despite there being steric strain present in the most substituted alkenes).


## Strongest Pi Bond



***Time Capsule:  
Zaitsev's rule follows  
this trend!!***

Cl, Br, I (not F)


### Addition of H-X to an Alkene



Summary:

Regiochemistry:

Stereochemistry:




For a reaction to take place, you need both:

Motive  $\rightarrow$  thermodynamic driving force


- \* 1) Make stronger bonds than you break
- 2) Formation of a weaker acid/base.
- 3) Entropy  $\rightarrow$  if you make a small gas molecule as a product.

Opportunity  $\rightarrow$  reasonable mechanism so that the activation energy of the reaction is not too high  
 $\Rightarrow$  movement of electrons




$$k = \text{reaction rate} = Ae^{-E_a/RT}$$

Increasing Temperature  $\rightarrow$  increases the number of molecules with enough energy to react



# Appendix 3



## Bond Dissociation Enthalpies

Bond dissociation enthalpy (BDE) is defined as the amount of energy required to break a bond homolytically into two radicals in the gas phase at 25°C.



| Bond   | $\Delta H^0$ | Bond  | $\Delta H^0$ | Bond  | $\Delta H^0$ |
|--|--------------|---|--------------|---|--------------|
| <b>H—H bonds</b>   |              | <b>C—C multiple bonds</b>                         |              | <b>C—Br bonds</b>                                 |              |
| H—H  | 435 (104)    | CH <sub>2</sub> =CH <sub>2</sub>                  | 727 (174)    | CH <sub>3</sub> —Br                               | 301 (72)     |
| D—D  | 444 (106)    | HC≡CH   | 966 (231)    | C <sub>2</sub> H <sub>5</sub> —Br                 | 301 (72)     |
| <b>X—X bonds</b>   |              | <b>C—H bonds</b>                                  |              | (CH <sub>3</sub> ) <sub>2</sub> CH—Br             | 309 (74)     |
| F—F  | 159 (38)     | CH <sub>3</sub> —H                                | 439 (105)    | (CH <sub>3</sub> ) <sub>3</sub> C—Br              | 305 (73)     |
| Cl—Cl  | 247 (59)     | C <sub>2</sub> H <sub>5</sub> —H                  | 422 (101)    | CH <sub>2</sub> =CHCH <sub>2</sub> —Br            | 247 (59)     |
| Br—Br  | 192 (46)     | (CH <sub>3</sub> ) <sub>2</sub> CH—H              | 414 (99)     | C <sub>6</sub> H <sub>5</sub> —Br                 | 351 (84)     |
| I—I  | 151 (36)     | (CH <sub>3</sub> ) <sub>3</sub> C—H               | 405 (97)     | C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> —Br | 263 (63)     |
| <b>H—X bonds</b>   |              | CH <sub>2</sub> =CH—H                             | 464 (111)    | <b>C—I bonds</b>                                  |              |
| H—F  | 568 (136)    | CH <sub>2</sub> =CHCH <sub>2</sub> —H             | 372 (89)     | CH <sub>3</sub> —I                                | 242 (58)     |
| H—Cl   | 431 (103)    | C <sub>6</sub> H <sub>5</sub> —H                  | 472 (113)    | C <sub>2</sub> H <sub>5</sub> —I                  | 238 (57)     |
| H—Br   | 368 (88)     | C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> —H  | 376 (90)     | (CH <sub>3</sub> ) <sub>2</sub> CH—I              | 238 (57)     |
| H—I  | 297 (71)     | HC≡C—H  | 556 (133)    | (CH <sub>3</sub> ) <sub>3</sub> C—I               | 234 (56)     |
| <b>O—H bonds</b>   |              | <b>C—F bonds</b>                                  |              | CH <sub>2</sub> =CHCH <sub>2</sub> —I             | 192 (46)     |
| HO—H   | 497 (119)    | CH <sub>3</sub> —F                                | 481 (115)    | C <sub>6</sub> H <sub>5</sub> —I                  | 280 (67)     |
| CH <sub>3</sub> O—H  | 439 (105)    | C <sub>2</sub> H <sub>5</sub> —F                  | 472 (113)    | C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> —I  | 213 (51)     |
| C <sub>6</sub> H <sub>5</sub> O—H                                    | 376 (90)     | (CH <sub>3</sub> ) <sub>2</sub> CH—F              | 464 (111)    | <b>C—N single bonds</b>                           |              |
| <b>O—O bonds</b>   |              | C <sub>6</sub> H <sub>5</sub> —F                  | 531 (127)    | CH <sub>3</sub> —NH <sub>2</sub>                  | 355 (85)     |
| HO—OH  | 213 (51)     | <b>C—Cl bonds</b>                                 |              | C <sub>6</sub> H <sub>5</sub> —NH <sub>2</sub>    | 435 (104)    |
| CH <sub>3</sub> O—OCH <sub>3</sub>                                   | 159 (38)     | CH <sub>3</sub> —Cl                               | 351 (84)     | <b>C—O single bonds</b>                           |              |
| (CH <sub>3</sub> ) <sub>3</sub> CO—OC(CH <sub>3</sub> ) <sub>3</sub> | 159 (38)     | C <sub>2</sub> H <sub>5</sub> —Cl                 | 355 (85)     | CH <sub>3</sub> —OH                               | 385 (92)     |
| <b>C—C single bonds</b>  |              | (CH <sub>3</sub> ) <sub>2</sub> CH—Cl             | 355 (85)     | C <sub>6</sub> H <sub>5</sub> —OH                 | 468 (112)    |
| CH <sub>3</sub> —CH <sub>3</sub>                                     | 378 (90)     | (CH <sub>3</sub> ) <sub>3</sub> C—Cl              | 355 (85)     |   |              |
| C <sub>2</sub> H <sub>5</sub> —CH <sub>3</sub>                       | 372 (89)     | CH <sub>2</sub> =CHCH <sub>2</sub> —Cl            | 288 (69)     |   |              |
| CH <sub>2</sub> =CH—CH <sub>3</sub>                                  | 422 (101)    | C <sub>6</sub> H <sub>5</sub> —Cl                 | 405 (97)     |   |              |
| CH <sub>2</sub> =CHCH <sub>2</sub> —CH <sub>3</sub>                  | 322 (77)     | C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> —Cl | 309 (74)     |   |              |
| C <sub>6</sub> H <sub>5</sub> —CH <sub>3</sub>                       | 435 (104)    |   |              |   |              |
| C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> —CH <sub>3</sub>       | 326 (78)     |   |              |   |              |

**Organic Chemistry is the study of carbon-containing molecules.**

**This class has two points.**

***The first point of the class is to understand the organic chemistry of living systems. We will teach you how to think about and understand the most amazing things on the planet!!***

Water is essential for life, you will learn why water has such special properties.

8/28/24

You will learn the secret structural reason proteins, the most important molecular machines in our bodies, can support the chemistry of life.

9/11/24

You will learn why when you take Advil for pain, exactly half of what you take works, and the other half does nothing.

9/25/24

You will learn how toothpaste works.

You will learn how a single chlorofluorocarbon refrigerant molecule released into the atmosphere can destroy many, many ozone molecules, leading to an enlargement of the ozone hole.

10/7/24

You will learn how medicines like Benadryl, Seldane, and Lipitor work.

You will learn how Naloxone is an antidote for an opioid overdose.

You will learn why Magic Johnson is still alive, decades after contracting HIV.

You will learn how MRI scans work.

***The second point of organic chemistry is the synthesis of complex molecules from simpler ones by making and breaking specific bonds.***

You will learn how to understand movies of reaction mechanisms like alkene hydration.

You will learn reactions that once begun, will continue reacting such that each product molecule created starts a new reaction until all the starting material is used up.

You will learn reactions that can make antifreeze from vodka.

You will learn a reaction that can make nail polish remover from rubbing alcohol.

You will learn how to look at a molecule and accurately predict which atoms will react to make new bonds, and which bonds will break during reactions.

You will learn how to analyze a complex molecule's structure so that you can predict ways to make it via multiple reactions starting with less complex starting molecules.

Mechanisms → movement of electrons and atoms in chemical reactions

1) Arrows in reaction mechanisms move electrons.

2) Arrows in reaction mechanisms, DO NOT move atoms

3) Arrows start at an

electron source → lone pair or a

and end at an  $\pi$  bond on an

electron sink species OR a


bond that

must break

↙  
an atom that can accept a new pair of electrons

4) Breaking a bond will occur to overfilling the valence shell of an atom.






a) → lone pair on N atom makes a new bond to H

b) → O-H sigma bond breaks to prevent overfilling the valence shell of H

Mistake to avoid → moving an atom with an arrow



WRONG!! → Move the

electrons with an arrow not an H atom!

**Nucleophile** → electron rich species that has an electron rich  $\pi$  bond or a lone pair that takes part in a bond forming step

↓

analogous to a Lewis base

Ex.



**Electrophile** → electron deficient species that contains an atom that serves as a sink for an arrow from a nucleophile


↓

analogous to a Lewis acid


OR

a molecule that has a weak bond

Ex. Full  $\oplus$



Partial  $\oplus$



Weak bond



# The 4 Most Important Mechanistic Elements

The following are expressed from the point of view of the carbon-containing molecule taking part in a reaction

1) **Make a bond** between a nucleophile and electrophile.

⇒ A nucleophile and electrophile are both present and a bond can be made.

2) **Break a bond** to give stable molecules or ions.

⇒ None of the other possibilities are likely and the fragments produced are relatively stable

3) **Add a proton**

⇒ Acid is present or the molecule is a strong base.

4) **Take a proton away**


⇒ Base is present or the molecule is a strong acid.

Notice → 1) is the reverse of 2) and 3) is the reverse of 4) .... and vice versa


# Mechanism Summary

The following questions and mechanistic elements are described from the point of view of the carbon-containing reagent, written in the form of a flowchart.

Is there a strong acid present or is the carbon-containing reagent a strong base?


**YES** 

**Add a proton**




**NO**

Is there a strong base present or is the carbon-containing reagent a strong acid?


**YES** 

**Take a proton away**




**NO**

Are there a nucleophile and electrophile present?


**YES** 

**Make a bond**




**NO**

Can a bond be broken to create stable molecules or ions?

**YES** 

**Break a bond**



**NO**

Think about alternative mechanistic elements








Addition of H-X to an Alkene

X = Cl, Br, I  
but not F



Summary:

Regiochemistry:

Stereochemistry:

Example:

