The Golden Rules of Organic Chemistry

Your goal should be to understand, not memorize, the material presented in your organic chemistry course. The following principles should be learned as you begin your study of organic chemistry, then used as a solid foundation for building your understanding throughout the course. These simple ideas explain a great deal about the structures and properties of organic molecules, as well as the characteristic ways in which they react. Thoroughly understanding the following three key principles and related ideas will allow you to develop an intuitive feel for organic chemistry that avoids the necessity of resorting to the far less effective use of extensive memorization.

A. Predicting Structure and Bonding

- 1. In most stable molecules, all the atoms will have filled valence shells.
- 2. Five- and six-membered rings are the most stable.
- 3. There are two possible arrangements of four different groups around a tetrahedral atom.

B. Predicting Stability and Properties

- 4. The most important question in organic chemistry is "Where are the electrons?"
- 5. Delocalization of charge over a larger area is stabilizing.
- 6. Delocalization of unpaired electron density over a larger area is stabilizing.
- 7. Delocalization of pi electron density over a larger area is stabilizing.

C. Predicting Reactions

- 8. Reactions will occur if the products are more stable than the reactants and the energy barrier is low enough.
- 9. Functional groups react the same in different molecules.
- 10. A reaction mechanism describes the sequence of steps occurring during a reaction.
- 11. Most bond-making steps in reaction mechanisms involve nucleophiles reacting with electrophiles.




FIGURE 21.2 The molecular orbital representation of the π bonding in benzene.




FIGURE 21.3 Orbitals for the π system of benzene. (a) Cartoon representations of the six calculated orbitals that chemists routinely draw. These pictures accentuate the fact that various combinations of parallel 2p orbitals lead to the π system of benzene. (b) Calculated orbitals. The three lowest in energy are occupied with electrons (see Figure 21.2). The lowest of these orbitals is the image most chemists use for the π system of benzene: a torus of electron density above and below the ring.

Hückel's Aromaticity Criteria


- 1)
- 2)
- 3)
- 4)




Benzene




Tropylium Ion




Cyclopentadienyl anion



Pyrrole



Pyridine



Annulene



Cyclobutadiene



Cyclooctatetraene

Naphtalene

Anthracene

 $\mathsf{Benzo}[\alpha]\mathsf{pyrene}$

Phenoxide anion

Benzyl cation

Benzyl radical

$$\mathbf{E}^{\oplus}$$

$$\mathbf{H}$$

$$\mathbf{H$$

Н

Reagents

Wicked strong electrophile

 \mathbf{E}^{\oplus}

 χ^{\oplus}

Halogenation X2, FeX3

$$X = Br, Cl$$


Nitration H₂SO₄/HNO₃

$\underline{Sulfonation}\;H_2SO_4/SO_3$

Reagents

Friedel-Crafts Alkylation R-X, AlX₃

 R^{\oplus}



Note this is a carbocation, so it will rearrange if it is a primary or a rearrangmentprone secondary cation

Friedel-Crafts Acylation RCOCl, AlCl₃

Other notes: 1) It is hard to stop the Friedel-Crafts alkylation after one alkyl group adds (because alkyl groups are "good", that is, activating), but it can be done. 2) Neither Friedel-Crafts reaction works if there is already an electron withdrawing (bad) group on the ring.

Arenium ion stabilizing interactions

A) Pi donation, a resonance effect for atoms with lone pairs attached to the ring

B) Hyperconjugation for alkyl groups attached to the ring

Arenium ion destabilizing interaction

A) Inductive effect of electronegative atoms or groups attached to the ring

	Strongly activating	−ÄH₂	— ÑHR	$-\ddot{\mathrm{N}}\mathrm{R}_{2}$	— <u>ё</u> н	−ÿR		
Ortho-Para Directing	Moderately activating	O — NHCR	-NHCAr	−öcr	−öcar			GOOD
Ortho-Pa	Weakly activating	—R	$\overline{}$					
	Weakly deactivating	- <u>;</u> :	-çı:	— <u>ÿ</u> ι:	— <u>;;</u> :			UGLY
Meta Directing	Moderately deactivating	—СН	O -CR	—сон 0	-cor	$-\text{CNH}_2$	-soh	-c≡n
Meta	Strongly deactivating	$-NO_2$	—NH ₃ +	—сғ ₃	—cci ₃			BAD

ORTHO

ORTHO

META

Η,

H[′]

PARA

