

NAME (Print): _____

SIGNATURE: _____

**Chemistry 320N
Dr. Brent Iverson
5th Homework
February 18, 2026**

**Please print the
first three letters
of your last name
in the three boxes**

--	--	--


For all of the following mechanism sheets, use **arrows to indicate movement of all electrons, write all lone pairs, all formal charges, and all the products for each step.** Remember, I said all the products for each step. **IF A NEW CHIRAL CENTER IS CREATED IN AN INTERMEDIATE, MARK IT WITH AN ASTERISK AND LABEL THE MOLECULE AS “RACEMIC” IF APPROPRIATE. There is no need to use wedges and dashes for intermediates. FOR ALL CHIRAL PRODUCTS YOU MUST DRAW ALL ENANTIOMERS WITH WEDGES AND DASHES AND WRITE “RACEMIC” IF APPROPRIATE.** Next to each set of equilibrium arrows in each mechanism, write which of the 4 most common mechanistic elements describes each step (make a bond, break a bond, etc.).

When predicting mechanisms, remember to:


1. **Identify which bonds will be made and broken** in overall reaction (provided).
2. **Avoid “mixed media errors”**
 - a. In acid, all intermediates are positively-charged or neutral
 - b. In base, all intermediates are negatively-charged or neutral
 - c. Only in neutral solutions might there be both positively-charged and negatively-charged intermediates.
3. **Proton transfers are fast** compared to other steps (**when in doubt transfer a proton**).
4. **Analyze** each intermediate to **PREDICT** the next step.


Here are the four mechanistic elements. The vast majority of mechanistic steps are one of these, and they are entirely predictable based on the personalities of the species involved:

1. **Make a Bond**
2. **Break a Bond**
3. **Add a Proton**
4. **Take a Proton Away**







Acid Catalyzed Anhydride Hydrolysis







Acid Catalyzed Ester Hydrolysis







Base-Promoted Ester Hydrolysis - Saponification









Acid Promoted Amide Hydrolysis






Acid Chlorides Reacting with Amines








Grignard Reacting with Esters




Chemist Opens Flask

2)








Reduction of Esters with LiAlH₄




Chemist Opens Flask







Acid Promoted Nitrile Hydrolysis



OK, the one on the next page has a twist! It has a unique step that you have not seen before. Hint: It has a lot to do with a Lewis acid-Lewis base complex that behaves differently than you might expect.



Reduction of Amides with LiAlH₄



Note: In this reaction the chemist opens the flask and adds water in a second step that quenches any excess LiAlH₄. Therefore, you need a second step to add water when using this reaction in synthesis even though it is not shown in the mechanism above.