π-Way Recap

3 atom "π-ways" we have seen



Conjugation ⟶ "π way" ⟶ 4 atoms or more

↳ More than one π bond that overlaps



Not conjugated:





Butadiene

λ_{max} = 217 nm



β-Carotene

λ_{max} = 455 nm, 483 nm



11-*cis*-Retinal

λ_{max} = 380 nm



Lycopene

λ_{max} = 443 nm, 471 nm, 502 nm



Luteine

λ_{max} = 445 nm, 474 nm

White → reflects all wavelengths of visible light
Black → absorbs all wavelengths of visible light



FIGURE 20.5 (a) Visible light color-wavelength correlation.

*** We "see" the wavelengths reflected minus the wavelengths absorbed ***



(b) Approximate color of substance (reflected light) if a single wavelength (i.e., the wavelength listed on the numerical scale of the *x*-axis) is absorbed.



(c) Complementary colors on a color wheel.

Colored arrows are complementary

Generation of heat, Most molecules



Flourescence - Rigid Molecules, Not uncommon



Phosphorescence - “Glow in the Dark”, Rare



How vision works, the final edition!



Butadiene

λ_{max} = 217 nm



β-Carotene

λ_{max} = 455 nm, 483 nm



11-*cis*-Retinal

λ_{max} = 380 nm



Lycopene

λ_{max} = 443 nm, 471 nm, 502 nm



Luteine

λ_{max} = 445 nm, 474 nm

How vision works



Binds to an -NH_2 group from the amino acid lysine in the protein opsin



Absorbing the photon puts an electron into an antibonding π orbital → weaken π bond, especially around $C_{11}=C_{12}$ so bond can rotate back to more stable trans geometry

Molecule resets

A photon of visible light is absorbed by the retinal, isomerizing the *cis* bond to *trans*, dramatically altering the shape of the molecule



This change in shape causes a nerve impulse to be sent to the brain

Generation of heat, Most molecules



Flourescence - Rigid Molecules, Not uncommon



Phosphorescence - "Glow in the Dark", Rare



Flourescence - Rigid Molecules, Not uncommon



Fluorescein



Rhodamine

Phosphorescence - “Glow in the Dark”, Rare



Bioluminescence - Fireflies, Deep Sea Creatures - Chemical Reactions



http://photobiology.info/Branchini2.html



FIGURE 20.5 (a) Visible light color-wavelength correlation.

(b) Approximate color of substance (reflected light) if a single wavelength (i.e., the wavelength listed on the numerical scale of the *x*-axis) is absorbed.



(c) Complementary colors on a color wheel.

Colored arrows are complementary

Vanta Black → The "blackest" material









Science **322**, 238 (2008)

Preview



Extraordinarily Stable!

~36 kcal/mol more stable than expected

This is A LOT

Benzene

Called "aromaticity"

"aromatic" molecule

Pericyclic Reactions → π bonds and σ bonds interchange

→ Happens because the transition state is super stable

"aromatic" character of transition state

Otto!

↓

Diels-Alder Reaction

..... bonds being broken

..... bonds forming



s-cis butadiene



‡

3 π bonds being made or used – all ring atoms are sp^2 hybridized to begin with

"dienophile"



The above reaction gives a poor yield and was used only to illustrate the process → there are many, many known examples of Diels-Alder reactions

The following is the only Diels-Alder reaction you are responsible for in this class



Highly reactive dienophile

Cyclopentadiene



Endo → the ester groups are underneath the structure